

Impact Factor:

ISRA (India) = 6.317
ISI (Dubai, UAE) = 1.582
GIF (Australia) = 0.564
JIF = 1.500

SIS (USA) = 0.912
PIHII (Russia) = 3.939
ESJI (KZ) = 9.035
SJIF (Morocco) = 7.184

ICV (Poland) = 6.630
PIF (India) = 1.940
IBI (India) = 4.260
OAJI (USA) = 0.350

SOI: [1.1/TAS](https://doi.org/10.15863/TAS) DOI: [10.15863/TAS](https://doi.org/10.15863/TAS)

International Scientific Journal Theoretical & Applied Science

p-ISSN: 2308-4944 (print) e-ISSN: 2409-0085 (online)

Year: 2021 Issue: 11 Volume: 103

Published: 05.11.2021 <http://T-Science.org>

QR – Issue



QR – Article



Sh. Sadullaeva

Tashkent University of Information Technologies named after Muhammad al-Khwarizmi
Prof. of "Multimedia Technologies" department
Tashkent, Uzbekistan
orif_sh@list.ru

Z. Fayzullaeva

Tashkent University of Information Technologies named after Muhammad al-Khwarizmi
assistant of "Basics of Computer Science" department
Tashkent, Uzbekistan
nigor1802@mail.ru

INVESTIGATION ON SELF-SIMILAR ANALYSIS OF THE PROBLEM BIOLOGICAL POPULATION KOLMOGOROV-FISHER TYPE SYSTEM

Abstract: In this work we considered a parabolic system of two quasilinear reaction-diffusion equations for a biological population problem of the Kolmogorov-Fisher type describes the process of a biological population in a nonlinear two-component medium. We studied the qualitative properties of the solution to Cauchy problem based on self-similar analysis and its numerical solutions using the methods of modern computer technologies, to study the methods of linearization to the convergence of the iterative process with further visualization.

Key words: Cauchy problem, quasilinear, reaction-diffusion, biological population, numerical solutions.

Language: English

Citation: Sadullaeva, S. H., & Fayzullaeva, Z. (2021). Investigation on self-similar analysis of the problem biological population kolmogorov-fisher type system. *ISJ Theoretical & Applied Science*, 11 (103), 319-322.

Soi: <http://s-o-i.org/1.1/TAS-11-103-20> **Doi:**  <https://dx.doi.org/10.15863/TAS.2021.11.103.20>

Scopus ASCC: 2600.

Introduction

We can consider a parabolic system of two quasilinear reaction-diffusion equations for a biological population problem of the Kolmogorov-Fisher type in the following domain

$$Q = \{(t, x) : 0 < t < \infty, x \in \mathbb{R}^2\}$$
$$\begin{cases} \frac{\partial u_1}{\partial t} = \frac{\partial}{\partial x} \left(D_1 u_1^{\sigma_1} \frac{\partial u_1}{\partial x} \right) + k_1(t) u_1 \cdot (-u_2^{\beta_1}) \\ \frac{\partial u_2}{\partial t} = \frac{\partial}{\partial x} \left(D_2 u_2^{\sigma_2} \frac{\partial u_2}{\partial x} \right) - k_2(t) u_2 \cdot (-u_1^{\beta_2}) \end{cases} \quad (1)$$

$$\begin{aligned} u_1|_{t=0} &= u_{10}(x) \\ u_2|_{t=0} &= u_{20}(x) \end{aligned} \quad (2)$$

It describes the process of a biological population in a nonlinear two-component medium,

the diffusion coefficient of which is equal to $D_1 u_1^{\sigma_1}$ and $D_2 u_2^{\sigma_2}$, $\sigma_1, \sigma_2, \beta_1, \beta_2$ - positive real numbers, $u_1 = u_1(t, x) \geq 0$, $u_2 = u_2(t, x) \geq 0$ - sought solutions.

The Cauchy problem and boundary value problems for system (1) in the one-dimensional and multidimensional cases have been studied by many authors.[1]

The purpose of this work is to study the qualitative properties of the solution to problem (1), (2) based on self-similar analysis and its numerical solutions using the methods of modern computer technologies. Also the article has a purpose to study the methods of linearization to the convergence of the iterative process with further visualization. The main estimations of the solutions and the resulting free boundary have been found, which makes it possible

Impact Factor:

ISRA (India) = 6.317	SIS (USA) = 0.912	ICV (Poland) = 6.630
ISI (Dubai, UAE) = 1.582	PIHII (Russia) = 3.939	PIF (India) = 1.940
GIF (Australia) = 0.564	ESJI (KZ) = 9.035	IBI (India) = 4.260
JIF = 1.500	SJIF (Morocco) = 7.184	OAJI (USA) = 0.350

to choose appropriate initial approximations [...] Each of them has their own counting systems.

Now we will start constructing a self-similar system of equations for (1) - (2). It is a simpler system of equations for research.

We construct a self-similar system of equations by the method of nonlinear splitting.

Instead of (1)

$$u_1(t, x) = e^{k_1 t} v_1(t, x),$$

$$u_2(t, x) = e^{k_2 t} v_2(t, x)$$

This will lead (1) to the form:

$$\begin{cases} \frac{\partial v_1}{\partial \tau} = \frac{\partial}{\partial x} \left(D_1 v_1^{\sigma_1} \frac{\partial v_1}{\partial x} \right) + k_1 e^{((\beta_1+1)k_1 - (\delta_1+1)k_2)t} v_1 v_2^{\beta_1}, \\ \frac{\partial v_2}{\partial \tau} = \frac{\partial}{\partial x} \left(D_2 v_2^{\sigma_2} \frac{\partial v_2}{\partial x} \right) + k_2(t) e^{((\beta_2+1)k_2 - (\delta_2+1)k_1)t} v_1^{\beta_2} v_2, \end{cases} \quad (3)$$

$$v_1|_{t=0} = v_{10}(x), \quad v_2|_{t=0} = v_{20}(x),$$

After choosing $\sigma_1 k_1 = \sigma_2 k_2$, we can achieve the following forms of equation systems.

$$\begin{cases} \frac{\partial v_1}{\partial \tau} = \frac{\partial}{\partial x} \left(D_1 v_1^{\sigma_1} \frac{\partial v_1}{\partial x} \right) - a_1 \tau^{b_1} v_1 v_2^{\beta_1}, \\ \frac{\partial v_2}{\partial \tau} = \frac{\partial}{\partial x} \left(D_2 v_2^{\sigma_2} \frac{\partial v_2}{\partial x} \right) - a_2 \tau^{b_2} v_1^{\beta_2} v_2, \end{cases} \quad (4)$$

Where $a_1 = (\delta_1 k_1)^{b_1}$, $a_2 = (\delta_2 k_2)^{b_2}$

$$b_1 = [(\beta_1+1)k_1 - (\sigma_1+1)k_2] / \sigma_1 k_1, \quad b_2 = (\beta_2+1)k_2 - (\sigma_2+1)k_1 / \sigma_2 k_2$$

In the following, we can write one of the ways of auto-model systems for equation systems (4). It is done like in the following:

$$\begin{cases} \frac{d\bar{v}_1}{d\tau} = -a_1 \tau^{b_1} v_1 v_2^{\beta_1}, \\ \frac{d\bar{v}_2}{d\tau} = -a_2 \tau^{b_2} v_1^{\beta_2} v_2, \end{cases}$$

It has a solution in the following:

$$\bar{v}_1(t) = c_1(\tau + T)^{\gamma_1}, \quad \bar{v}_2(t) = c_2(\tau + T)^{\gamma_2}, \quad T > 0$$

And then systems of solution are sought in the following steps. (3)-(4)

$$v_1(t, x) = \bar{v}_1(t) w_1(\tau, x),$$

$$v_2(t, x) = \bar{v}_2(t) w_2(\tau, x),$$

here $\tau = \tau(t)$ is chosen in the following way:

$$\tau(t) = \int \bar{v}_1^{-\sigma_1}(t) dt = \frac{1}{\gamma_1 \sigma_1 + 1} (T + t)^{\sigma_1 \gamma_1 + 1}, \quad \gamma_1 \sigma_1 + 1 \neq 0$$

$$u \tau(t) = \ln(T + t), \quad \gamma_1 \sigma_1 + 1 = 0$$

Here we can choose equation system for $w_i(\tau, x)$, $i = 1, 2$

$$\begin{cases} \frac{\partial w_1}{\partial \tau} = \frac{\partial}{\partial x} \left(D_1 w_1^{\sigma_1} \frac{\partial w_1}{\partial x} \right) - \theta_1 (w_1 w_2^{\beta_1} - w_1) \\ \frac{\partial w_2}{\partial \tau} = \frac{\partial}{\partial x} \left(D_2 w_2^{\sigma_2} \frac{\partial w_2}{\partial x} \right) + \theta_2 (w_2 w_1^{\beta_1} - w_2) \end{cases},$$

$$\eta_1 = b_1 + 1 + \beta_1 (b_2 + 1) \neq 0$$

$$\eta_2 = -\beta_2 (b_1 + 1) + (b_2 + 1) \neq 0$$

It is as $\gamma_1 \sigma_1 > 0$, $\gamma_1 \sigma_1 = \gamma_2 \sigma_2$, $d_i > 0$. In this case we can rely on the $w_i(\tau(t), x) = y_i(\xi)$, $\xi = |x| / \tau_1^{1/2}$, $i = 1, 2$,

$$\begin{cases} \xi^{1-N} \frac{d}{d\xi} \left(\xi^{N-1} y_1^{\sigma_1} \frac{dy_1}{d\xi} \right) + \frac{\xi}{2\theta_1} \frac{dy_1}{d\xi} - \mu_1 (y_1 - y_1 y_2^{\beta_1}) = 0 \\ \xi^{1-N} \frac{d}{d\xi} \left(\xi^{N-1} y_2^{\sigma_2} \frac{dy_2}{d\xi} \right) + \frac{\xi}{2\theta_2} \frac{dy_2}{d\xi} + \mu_2 (y_2 - y_2 y_1^{\beta_2}) = 0 \end{cases} \quad (5)$$

Here $\mu_i = \frac{1}{\theta_i \sigma_i}$, $\theta_i = \begin{cases} 1, \\ \gamma_1^{-\sigma_1} \gamma_2^{\sigma_2}, \end{cases}$

The study of the qualitative properties of the system (1) - (2) made it possible to perform a numerical experiment depending on the values

We can choose the following equation system considering the fact that equation for $w_i(\tau, x)$ without little members is always auto-model.

included in the system of numerical parameters. For this purpose, the constructed asymptotic solutions were used as an initial approximation. In the numerical solution of the problem for the linearization of system (1) - (2), linearizations by the Newton and Picard methods were used. The method

Impact Factor:

ISRA (India)	= 6.317	SIS (USA)	= 0.912	ICV (Poland)	= 6.630
ISI (Dubai, UAE)	= 1.582	PIHII (Russia)	= 3.939	PIF (India)	= 1.940
GIF (Australia)	= 0.564	ESJI (KZ)	= 9.035	IBI (India)	= 4.260
JIF	= 1.500	SJIF (Morocco)	= 7.184	OAJI (USA)	= 0.350

of nonlinear splitting is proposed to solve the problem of a biological population.

$$\omega_h = \{x_i = ih, h > 0, i = 0, 1, \dots, n, hn = l\},$$

Temporary grid

$$\omega_{h_1} = \{t_j = jh_1, h_1 > 0, j = 0, 1, \dots, n, \tau m = T\}.$$

The main problem in nonlinear problems is the appropriate choice of the initial approximation and the way to linearize equation (3).

We replace problem (3) - (4) with an implicit difference scheme and obtain a difference problem with an error $O(h^2 + h_1)$.

$$\psi_1(t) = \bar{v}_1(t), \quad v_{10}(t, x) = \psi_1(t) \cdot \left(a - \frac{\sigma_1}{4} \xi^2\right)_+^{1/\sigma_1},$$

$$\psi_2(t) = \bar{v}_2(t), \quad v_{20}(t, x) = \psi_2(t) \cdot \left(a - \frac{\sigma_2}{4} \xi^2\right)_+^{1/\sigma_2},$$

$$\xi = \frac{x}{[\tau(t)]^{1/2}}, \quad \tau(t) = \int_0^t [\psi(y)] dy$$

$$(a)_+ \text{ means } (a)_+ = \max(0, a).$$

As a conclusion, the results of numerical experiments have shown the effectiveness of the proposed approach. Asymptotes of various solutions of the system of type (1) - (2) made it possible to simulate the processes of mutual reaction-diffusion in the form of visualization with animation.

All in all, we can emphasize the importance of a joint study of migration and demographic processes. To analyze the population dynamics of interacting populations, it is important to jointly study the processes of fertility, mortality, trophic interactions, and various migrations. The introduction of nonlinearity into migration flows is the first step towards an adequate description of spatio-temporal population dynamics.

References:

- Martynenko, A.V., & Tedeev, A.F. (2007). The Cauchy problem for a quasilinear parabolic equation with a source and no homogeneous density. *Comput. Math. Math. Phys.* 47, no. 2, pp. 238–248.
- Aripov, M., & Sadullaeva, Sh. (2013). To properties of the equation of reaction diffusion with double nonlinearity and distributed parameters. *Journal of Siberian Federal University. Mathematics & Physics*, Vol. 6, Issue 2, pp. 157–167.
- Sadullaeva, Sh. (2016). Numerical Investigation of Solutions to a Reaction-diffusion System with Variable Density. *Journal of Siberian Federal University. Mathematics & Physics*, 9(1), pp. 90-101.
- Sadullaeva, S.A., Khaydarov, A.T., & Kabilianova, F.A. (2019). *Modeling of Multidimensional Problems in Nonlinear Heat Conductivity in Non-Divergence Case*, 3rd International Symposium on Multidisciplinary Studies and Innovative Technologies, ISMSIT 2019 – Proceedings, 8932954.
- Sadullaeva, S.A., & Khojimurodova, M.B. (2018). Properties of solutions of the Cauchy problem for degenerate nonlinear cross systems with convective transfer and absorption, *Springer Proceedings in Mathematics and Statistics*, 264, pp. 183-190.
- Aripov, M., & Sadullaeva, S. (2015). An asymptotic analysis of a self-similar solution for the double nonlinear reaction-diffusion system, *J. Nanosyst. Phys. Chem. Math.*, 6 (6), pp. 793-802.
- Aripov, M., & Sadullaeva, S. (2015). Qualitative properties of solutions of a doubly nonlinear reaction-diffusion system with a source. *J. Appl. Math. Phys.*, 3, pp. 1090-1099.
- Muhamediyeva, D.K. (2019). The property of the problem of reaction diffusion with double nonlinearity at the given initial conditions. *International Journal of Mechanical and Production Engineering Research and Development*, Volume 9, Issue 3, IJMPERDJUN2019117, Pages 1095-1106.
- Mersaid, A. (2018). The Fujita and secondary type critical exponents in nonlinear parabolic equations and systems. *Springer Proceedings in Mathematics and Statistics*, 268, pp. 9-23.
- Muhamediyeva, D.K. (2018). Properties of self-similar solutions of reaction-diffusion systems of quasilinear equations. *International Journal of Mechanical and Production Engineering Research and Development*, 8 (2), pp. 555-566.
- Sadullaeva, S.A., & Paradaeva, G. (2018). Numerical Investigation of a System Reaction-Diffusion with Double Nonlinearity. *Journal of Mathematics, Mechanics and Computer Science*, 86 (3), 58-62.

Impact Factor:

ISRA (India) = 6.317
ISI (Dubai, UAE) = 1.582
GIF (Australia) = 0.564
JIF = 1.500

SIS (USA) = 0.912
PIHIQ (Russia) = 3.939
ESJI (KZ) = 9.035
SJIF (Morocco) = 7.184

ICV (Poland) = 6.630
PIF (India) = 1.940
IBI (India) = 4.260
OAJI (USA) = 0.350

12. Aripov, M., Djabbarov, O., & Sadullaeva, Sh. (2021). *Mathematic modeling of processes describing by double nonlinear parabolic equation with convective transfer and damping*. AIP Conference Proceedings 2365, 060008 (2021); <https://doi.org/10.1063/5.0057492>
13. Aripov, M., Sadullaeva, S., & Iskhakova, N. (2020). *Numerical Modeling Wave Type Structures in Nonlinear Diffusion Medium with Dumping*. 4th International Symposium on Multidisciplinary Studies and Innovative Technologies, ISMSIT 2020 - Proceedings this link is disabled, 2020, 9254988
14. Sadullaeva, S., Fayzullaeva, Z., & Nazirova, D. (2020). *Numerical Analysis of Doubly Nonlinear Reaction-Diffusion System with Distributed Parameters*. 4th International Symposium on Multidisciplinary Studies and Innovative Technologies, ISMSIT 2020 - Proceedings this link is disabled, 2020, 9255106.
15. Nematov, A., Nazirova, E.S., & Sadikov, R.T. (2021). On numerical method for modeling oil filtration problems in piecewise-inhomogeneous porous medium, IOP Conference Series: *Materials Science and Engineering* this link is disabled, 1032(1), 012018
16. Nazirova, E., Nematov, A., Sadikov, R., & Nabiyeu, I. (2021). *One-Dimensional Mathematical Model and a Numerical Solution Accounting Sedimentation of Clay Particles in Process of Oil Filtering in Porous Medium*. Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics) this link is disabled, 2021, 12615 LNCS, pp. 353–360.
17. Ravshanov, N., Nazirova, E.S., & Pitolin, V.M. (2019). Numerical modelling of the liquid filtering process in a porous environment including the mobile boundary of the oil-water section. *Journal of Physics: Conference Series* this link is disabled, 1399(2), 022021.
18. Anarova, S., & Ismoilov, S. (2021). *Nonlinear mathematical model of stress-deformed state of spatially loaded rods with account for temperature*. AIP Conference Proceedings this link is disabled, 2021, 2365, 070019.
19. Anarova, S.A., Ismoilov, S.M., & Abdirozиков, O.S. (2021). *Software of Linear and Geometrically Non-linear Problems Solution Under Spatial Loading of Rods of Complex Configuration*, Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics) this link is disabled, 2021, 12615 LNCS, pp. 380–389.